

# 耳多贊邦點耳液劑

## Otozambon Ear Drops

衛署藥輸字第023964號



成份：Each 1ml contains:

Polymyxin B sulphate .....	10,000 I.U.
Neomycin sulphate .....	5 mg
(equivalent to Neomycin basis 3.75mg)	
Lidocaine hydrochloride .....	40 mg
Benzalkonium chloride .....	0.1 mg
Propylene glycol .....	0.44 ml
Glycerol .....	0.265 ml
Purified water to .....	1 ml

特性：OTOZAMBON 適於耳科感染症之治療。此劑含有對變形菌及 Pyocianeous 等有效之廣汎抗生素 Polymyxin B sulfate 及 Neomycin 外，含有麻醉劑 Lidocaine 解除耳炎時常見之疼痛現象。因由數種 抗菌劑之協和作用 OTOZAMBON 能消炎、解除疼痛及膿。

適應症：細菌性引起之耳部感染及耳部外聽道發炎。

用法及用量：本藥須由醫師處方使用。

成人：一次滴於耳孔內4~5滴，一日2~4次。

小孩：一次滴於耳孔內2~3滴，一日3~4次。

包裝：每支8公撮瓶裝。

禁忌：鼓膜有穿孔之場合對所含成份有過敏性之患者。

一般注意：

- (1) 長期使用抗生素場合會致抗藥性之真菌及其他微生物之過度繁殖，所以有這種場合須採取適當措施。
- (2) 對鼓膜有穿孔或已經過長期之中耳炎患者須慎重調節使用。
- (3) 不要繼續10日以上。
- (4) 與 Kanamycin, Paromomycin, Streptomycin 及 Gentamycin 合用時可能會引起過敏，醫師要留意此點。

副作用：要與留意此劑所含 Neomycin 之會引起過敏現象之可能性。

保存：25°C 以下避光儲存。

製造廠：Zambon SPA

Office: Via Lillo del Duca n. 10-12, 20091 Bresso Milan Italy

Plant: Via Della Chimica, 9 36100 Vicenza, Italy

藥商：幸生實業股份有限公司

地址：台北市中山區松江路520號6樓

電話：(02)2596-9138

# Otozambon®

---

## COMPOSITION

Polymyxin B sulfate 1,000,000 IU - Neomycin 0.375 g (as Neomycin sulfate) - Lidocaine hydrochloride 4 g - Benzalkonium chloride 0.01 g - Water, propylene glycol and glycerol q.s.p. 100 ml.

## PROPERTIES

OTOZAMBON® is a product particularly indicated in the treatment of several ear affections.

Besides Neomycin and Polymyxin B, broad antibacterial spectrum antibiotics, the elective activity of which respectively on *Proteus* and *Pyocianus* is well-known, the product contains Lidocaine, which, due to its anesthetic properties, relieves the painful symptoms accompanying otologic infections.

Because of the synergic action of these antibacterial agents, OTOZAMBON® provides a useful means in the treatment of infectious forms and furthermore allows to hasten the disappearance of phlogistic signs and mucopurulent secretions, to relieve the painful or pruriginous subjective symptomatology.

## INDICATIONS

Acute and chronic otitis externa. Acute otitis media in the pre- and post-perforative stage and chronic secretive otitis. Suppurative complications following ear and mastoid surgery (radial mastoidectomy, plastic surgery on the tympan, atticotomy, fenestration, etc.).

## USE AND DOSAGE

OTOZAMBON® must be instilled with the proper dropper in the external auricular tube at the dose for adults of 4-5 drops, 2-4 times daily according to the seriousness of the morbid form. In children the recommended dose is of 2-3 drops, 3-4 times daily. The treatment period varies in relation to the promptness of the therapeutic result.

The high tolerability of the preparation allows, if necessary, also prolonged treatment.

## PACKING

8 ml bottle with dropper.

**Zambon** SPA  
VICENZA (ITALY)